



Plaquenil and Your Eye Health

What is Plaquenil?

Plaquenil® (Hydroxychloroquine sulfate) is a drug used to treat certain **autoimmune diseases**. Some diseases treated with Plaquenil include:

- **Lupus**, which causes fever, rashes, skin problems, and other symptoms.
- **Rheumatoid arthritis**, a condition that causes pain and swelling in the joints of your hands and feet.
- **Sjögren's syndrome**, which causes dry eyes and dry mouth.

Plaquenil lowers your immune system's ability to cause **inflammation**. This can help control symptoms like rashes, skin and mouth sores and joint pain.

Eye Words to Know

Autoimmune disease: When your immune system can't tell the difference between healthy cells and threatening germs, like viruses or bacteria. Your body then attacks its own healthy tissues.

Inflammation: When part of the body becomes red, irritated, swollen, or painful. This can happen when the body is fighting an infection or in response to an injury.

Retina: Layer of cells lining the back wall inside the eye. This layer senses light and sends signals to the brain so you can see.

Plaquenil and your eyes

A rare side effect of Plaquenil is damage to the eye's **retina**. Using Plaquenil for a long period of time may harm the retina, causing serious vision loss.

People with retinal damage from Plaquenil are not aware at first that they are losing vision. Unfortunately, once they lose a severe amount of vision loss, it is permanent. If you take Plaquenil, it is very important to see an ophthalmologist regularly. Your ophthalmologist will check your retina for problems before serious damage occurs.

Who is at risk for vision loss with Plaquenil?

Some people who take Plaquenil have a higher risk of retinal damage than others. They include:


- people taking Plaquenil for 5 or more years
- people taking Plaquenil at higher doses than the maximum recommended daily dose of 5.0 mg/kg real weight
- people who already have retinal disease
- people who already have kidney or liver disease
- people over 60 years old
- people who lose a lot of weight while taking Plaquenil. Tell your doctor if this happens as your dose may need to be lowered.

When should you see an ophthalmologist when taking Plaquenil?

If you take Plaquenil, here is when to see an ophthalmologist:

- **Before Plaquenil treatment begins.** You will have a “baseline” eye exam. This measures the health of your eyes and looks for retinal or macular disease.
- **Once a year while taking Plaquenil.**
- **After five years of Plaquenil treatment,** for those on acceptable doses without major risk factors, see your ophthalmologist annually (once per year).

For those on acceptable doses with major risk factor(s) for retinal damage from Plaquenil, you may need to see your ophthalmologist more often. Special tests can help find early damage to the retina before serious problems develop.



If you take Plaquenil, it is important to see an ophthalmologist while you take this medicine.

How do ophthalmologists look for Plaquenil damage?

Your ophthalmologist may use these tests to look for retina damage from Plaquenil:

- **Visual field test:** During this test you sit down and look into a bowl-like machine. Small lights flash at different points inside the bowl. When you see the lights, you press a button. This test can help find if blind spots have developed in your field of vision.
- **OCT imaging:** This test makes a detailed, three-dimensional image of your eye. Your ophthalmologist can use this image to look for early retinal damage.
- **Multifocal ERG:** This test looks at cells in the retina called rods and cones. The test measures how well these cells respond to light. It can find retinal damage from Plaquenil.
- **Photos:** A special camera takes pictures of the retina. Photos show damaged areas as small spots of light.

What happens if retinal damage is found?

If your ophthalmologist finds any signs of retinal damage, he or she will tell your primary care doctor or inflammatory disease specialist to stop Plaquenil treatment immediately. This will help you avoid central vision loss. Unfortunately, vision loss from Plaquenil is not reversible, and there is no known therapy. So early detection of early retinal changes and stopping Plaquenil are the only steps that will protect your vision. Your other doctor will find another treatment for your disease.

Summary

Plaquenil is a drug used to control symptoms from certain immune system diseases. If you take Plaquenil, it is important to see an ophthalmologist before and while you take this drug. This is because Plaquenil can cause damage to your eye's retina.

Your ophthalmologist will check the health of your retina. If he or she finds damage from Plaquenil, you will need to stop taking this drug. Your primary care doctor or inflammatory disease specialist will find a different treatment for you.

If you have any questions about your eyes or your vision, talk to your ophthalmologist. He or she is committed to protecting your sight.

Get more information about eye health from EyeSmart—provided by the American Academy of Ophthalmology—at aao.org/eyesmart.

COMPLIMENTS OF:

LONGWOOD EYE & LASIK

180 Daggett Drive
33 Riddell Street
354 Main Street
33 Electric Avenue
800.676.5050

West Springfield
Greenfield
Gardner
Fitchburg
www.longwoodeye.com